





Captan 800 WG

FEATURES & BENEFITS:

- No foaming
- No unpleasant smell
- Excellent mixing and dispersing characteristics
- Minimal particle size = ensures excellent coverage
- Available in 10kg (paper) bags with plastic liner



PARTICLE SIZE ANALYSIS:



Melpat International
4/22 Parry Ave, Bateman, WA 6150

Phone: (08) 9312 3200; Fax: (08) 9312 3233

www.melpat.com.au



DIRECTIONS FOR USE - EXTRACT

Please note: this is only an extract from the “Directions for Use” Table. For full details, please refer to the complete Label.

RESTRAINTS:

DO NOT apply this product by aircraft.

Crop	Disease	State	Rate /100 litre water	Critical Comments	
RATE In the following table all rates given are for dilute spraying. For concentrate spraying, refer to the Application section in the GENERAL INSTRUCTIONS.				For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether apply this product by dilute or concentrate spraying methods. DO NOT use at concentrate rates greater than 250g/ 100L water.	
Apples	Black Spot (<i>Venturia inaequalis</i>)	All States	Dilute Spraying 125g/ 100L Concentrate spraying Refer to Mixing/ Application section of label	Apply from pink stage onwards following a copper spray at green tip, then at 7 day intervals until petal fall. After petal fall apply as a cover spray at 10-14 day intervals. The risk of russet may be increased in some varieties. Where this is of concern use alternative materials between full bloom and second cover. Apply to point of run-off. Apply no more than five applications per season.	
Pears	Black Spot (<i>Venturia pirina</i>)				
Grapes	Black Spot				Apply at 50% bud-burst and repeat application 10-14 days later. Apply to point of run-off. Apply no more than five applications per season.
	Grey Mould				Apply in a program of sprays applied at early flowering, 80-100% cap fall, pre-bunch closure, veraison and pre-harvest. Apply to point of run-off. Apply no more than five applications per season.
	Downy Mildew				Apply from flowering onwards, at intervals of 10-14 days. Apply to point of run-off. Apply no more than five applications per season.
	Phomopsis Cane and Leaf Blight				Commence application at early bud-burst and repeat at 7 to 14 day intervals until prior to flowering to maintain a cover of new grape foliage. Apply no more than 5 applications per season. Use the shorter application interval where there is a rapid shoot growth and conditions are favourable for phomopsis infection (wet weather). Adjust spray application equipment to obtain a thorough coverage of grape buds and shoots.
Stone fruit (except apricots)	Blossom Blight & Brown Rot				Apply at any of the following times as part of a spray program to control blossom blight and brown rot– pink bud, 10% blossom, full bloom, petal fall and shuck fall and pre-harvest sprays at 6, 3 and 1 week prior to harvest. Apply to point of run-off. Apply no more than five applications per season. DO NOT apply to apricots as damage to fruit and foliage may occur.
Strawberries	Grey Mould	125g/100L	Apply every 10 days commencing at blossom stage. Apply to point of run-off.		
	Phytophthora Fruit Rot Black Spot Scorch Leaf Blight	200g	Apply no more than five applications per season		

WITHHOLDING PERIODS:

STRAWBERRIES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

APPLES, PEARS, GRAPES, STONE FRUIT: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

DO NOT USE TREATED PRODUCE FOR STOCK FOOD WITHIN 7 DAYS OF TREATMENT

Melpat International
4/22 Parry Ave, Bateman, WA 6150

Phone: (08) 9312 3200; Fax: (08) 9312 3233

www.melpat.com.au

